

Fertilgold® SOLUBLE POWDER

INSTRUCTIONS

- FERTILGOLD® SOLUBLE POWDER should be applied when tissue and soil analysis or experience identifies the need and proper rate of application for this product.
- Adds naturally occurring materials containing humic and fulvic acids to soil that enhances plant micronutrient availability and uptake.
- Best results will be obtained when application is concentrated in the active root zone.
- Can be applied in combination with compatible plant growth regulators, pesticides, or other granular fertilizer blends or mixes. If compatibility is in question, jar test a small quantity.
- Apply with sufficient water to ensure uniform coverage.
- FERTILGOLD® SOLUBLE POWDER can be applied to all soil types for all crops.
- Application timing, intervals, and rates may vary according to individual crop requirement, available nutrient levels in the soil.
- Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation.
- Consult your local FERTILGOLD® Representative or other agricultural specialist for crop-specific recommendations.

This product is intended as a supplement to a complete fertilizer program and will not, by itself, provide the nutrients normally required by agricultural crops.

STORAGE AND DISPOSAL

Keep product in original bag. Do not transfer into food or drink containers. Always dispose of bag in accordance with local, state, and/or federal regulations.

0-0-0 GUARANTEED ANALYSIS

Also contains Non-Plant Food Ingredients:

70.0% - 80.0%+ Total Humic and Fulvic Acids

Derived from Oxidized Leonardite

*Determined using V&B (Verploegh and Brandvoid) Method

Designed to be applied to the soil.

Applications can be made as often as every 15 - 30 days as needed.

APPLICATION

CROP TYPE	APPLICATION RATE
Vegetable crops, Strawberries	Up to 2 lb/ac (2.25 kg/ha), diluted with 25 gal (100 liters) of water At planting: Banded, side dressed or injected in furrow At flower set and again 2 weeks prior to harvest: with irrigation
Row crops	Up to 3 lb/ac (3.36 kg/ha), diluted with 25 gal (100 liters) of water At planting: Banded, side dressed or injected in furrow At flower set and again 2 weeks prior to harvest: with irrigation
Citrus, Olives, Deciduous Fruit & Nut Trees	Up to 4 lb/ac (4.5 kg/ha), diluted with 25 gal (100 liters) of water At flower set and again 2 weeks prior to harvest: with irrigation

SAFETY

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

A brown to black powder and dust having a mild characteristic odor. Ingestion of this product may cause gastrointestinal irritation or pain.

CONDITIONS OF SALE

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions, and other factors are beyond the control of the seller.

Net Wt. 25 kg
Net Wt. 55 lbs

© 2017–2020 *Registered Trademarks of and
Produced by: Bio Huma Netics, Inc.
1331 W. Houston Ave., Gilbert, AZ 85233
480.961.1220
FGO-648-03162020

Fertilgold.com

