

## INSTRUCTIONS

- FERTILGOLD® K HUME® should be applied when tissue and soil analysis identifies the need and proper rate of application for this product.
- Adds naturally occurring materials containing humic and fulvic acids activated with an alkaline potassium that becomes an excellent soil conditioner.
- Best results will be obtained when application is concentrated in the active root zone.
- Can be applied in combination with compatible granular fertilizers. Avoid spreading during high humidity, or mixing with high moisture fertilizers such as urea.
- Apply FERTILGOLD® K HUME® by broadcast whether through air or ground machine. FERTILGOLD® K HUME® can be applied to all soil types for agriculture fields, ornamentals, lawns, gardens, and landscapes.
- Application timing, intervals, and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop.
- Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation.
- Consult your local FERTILGOLD® Representative or other agricultural specialist for crop-specific recommendations.

This product is intended as a supplement to a complete fertilizer program and will not, by itself, provide all of the nutrients normally required by agricultural crops.

### STORAGE AND DISPOSAL

Keep product in original bag. Do not transfer into food or drink containers. Always dispose of bag in accordance with local, state, and/or federal regulations.

0-0-3

## GUARANTEED ANALYSIS

Soluble Potash (K<sub>2</sub>O) ..... 3.0%

Derived from: Potassium Hydroxide

### ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

Total Humic Acid (HA) and  
Fulvic Acid (FA) ..... 60.0%-70.0%  
(Determined using Colorimetric Test Method)

### Other Test Methods:

Total Humic Acids (HA) ..... 40.0%-45.0%  
(Analysis using the HPTA or ISO 19822 Test Method)

Total Humic Acids (HA) ..... 35.0%-40.0%  
(Analysis using the CDFA Test Method)

Derived from: Oxidized Leonardite

## SOIL AMENDMENT

Applications can be made as often as  
every 30 days as needed.

## APPLICATION

METHOD	SUGGESTED RATE	
	Field Crops	Tree or Vine Crops
Soil banded or side dressed	Up to 75 lb/acre, 75 kg/hectare	Up to 150 lb/acre, 150 kg/hectare
Soil broadcast incorporated	Up to 150 lb/acre, 150 kg/hectare	Up to 300 lb/acre, 300 kg/hectare

## SAFETY

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

*A dark reddish brown to black particles/dust having no odor. Ingestion of this product may cause gastrointestinal irritation or pain.*

CLASSIFICATION: HAZARD CATEGORY 3 - MILD SKIN IRRITATION

SIGNAL WORD: WARNING

HAZARD STATEMENT: H316 - WARNING – causes mild skin irritation

PRECAUTIONARY STATEMENT: P332+P313; If skin irritation occurs: Get medical attention/advice.

### CONDITIONS OF SALE

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions, and other factors are beyond the control of the seller.

Approved for  
organic  
production.



For Organic Use

Net Wt. 50 lb  
Net Wt. 22.68 kg