LIQUID HA 15

INSTRUCTIONS

- FERTILGOLD®* LIQUID HA 15 should be applied when tissue and soil analysis identifies the need and proper rate of application for this product.
- Adds naturally occurring materials containing humic acids to soil that may enhance plant micronutrient availability and uptake.
- Best results will be obtained when application is concentrated in the active root zone or when applied directly to the soil followed by a shallow cultivation.
- Can be applied alone or with liquid fertilizers with a pH of 7.0 or higher. If compatibility is in question, jar test a small quantity.
- FERTILGOLD® LIQUID HA 15 can be applied to all crops including fruits, vegetables, trees, vines, flowers, ornamentals, turf, grains, and other row crops.
- Avoid spreading on live crops, during high temperatures, or in times of drought.
- Application timing, intervals, and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop.
- Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation.
- Consult your local FERTILGOLD® Representative or other agricultural specialist for crop-specific recommendations.

This product is intended as a supplement to a complete fertilizer program and will not, by itself, provide the nutrients normally required by agricultural crops.

STORAGE AND DISPOSAL

FGO-000-01112023

Keep product in original container. Do not transfer into food or drink containers. Triple rinse when empty for recycling. Always dispose of container in accordance with local, state, and/or federal regulations. Do not store this product below 50°F (10°C) or above 90°F (30°C).

© 2017–2023 *Registered Trademarks of and Produced and Guaranteed by: Bio Huma Netics, Inc. 1331 W. Houston Ave., Gilbert, AZ 85233 480.961.1220



0-0-3.5 GUARANTEED ANALYSIS

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

Total Humic Acid (HA) and

Fulvic Acid (FA) 15.0%

(Determined using Colorimetric Test Method)

Derived from: Oxidized Leonardite

Other Test Method:

SHAKE WELL BEFORE USE.

Applications can be made as often as every 15 to 30 days as needed.

APPLICATION

METHOD	SUGGESTED RATE	
	Field Crops	Tree or Vine Crops
Soil banded, injected, side dress, drip tape, or micro sprinklers	up to 1 gallon/ac up to 10 liters/ha	up to 2 gallons/ac up to 20 liters/ha
Sprinklers: solid set, drag lines, linear, or pivot @ 100% speed	up to 2 gallons/ac up to 20 liters/ha	up to 4 gallons/ac up to 40 liters/ha
Soil broadcast spray incorporated, flood or furrow irrigated	up to 4 gallons/ac up to 40 liters/ha	up to 8 gallons/ac up to 80 liters/ha

pH: 9.0 - 10.0

9.09 lb/gal @ 68°F 1.09 kg/L @ 20°C

LIQUID HA 15

SAFETY

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.html

A brown liquid having a mild odor.

CONDITIONS OF SALE

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions, and other factors are beyond the control of the seller.



2.5 gallons • Net Wt. 22.73 lb 9.46 liters • Net Wt. 10.31 kg

FERTILGOLD.COM

LIQUID HA 15